

SUPREP – Colonoscopy Prep Instructions – PLEASE READ CAREFULLY

DON'T FORGET
TO FILL YOUR PRESCRIPTION FOR

SUPREP
AT YOUR PHARMACY

IT IS VERY IMPORTANT FOR YOU TO CAREFULLY
FOLLOW THESE INSTRUCTIONS.

**A COMPLETE COLONOSCOPY CANNOT BE
PERFORMED ON AN UNCLEAN COLON!**

BLOOD THINNERS

- 5 Days prior to your exam stop taking Plavix (Clopidogrel) and Coumadin (Warfarin), Effient (Prasugrel)
- 2 Days prior to your exam stop taking Pradax (Dabigatran), Xarelto (Rivaroxaban), and Eliquis (Apixaban)
- Please check with your cardiologist about stopping these medications

5 DAYS PRIOR TO YOUR EXAM:

- If you have diabetes, your primary care provider will advise you how to manage your medications prior to the colonoscopy (e.g. insulin).

3 DAYS PRIOR TO YOUR EXAM:

- Avoid nuts, seeds, whole wheat bread, beans, corn, raw vegetables or fruits with seeds or skin, as they may adversely influence the quality of your exam.

2 DAYS PRIOR TO YOUR EXAM:

- You may eat a regular dinner, but stop eating solid foods after 7:00 pm.
- You may continue to drink clear liquids. This means liquids you can see through such as apple juice, water, Sprite, Jell-O, coffee (no creamer), tea, clear chicken or beef broth, clear Gatorade/Powerade, or Crystal Light. No milk or pulp products. Avoid anything with red dye.

1 DAY PRIOR TO YOUR EXAM:

- **Restrict your diet to clear liquids. No solid food until after the procedure. Stay hydrated!**
- At 5:00 PM, complete steps 1 through 4 using one 6-ounce Suprep bottle.
 - Step 1: Pour one 6 ounce bottle of Suprep liquid into the mixing container
 - Step 2: Add cool drinking water to the 16 ounces of plain water
 - Step 3: Drink all the liquid in the container
 - Step 4: Over the next hour drink 2 additional 16 ounces of plain water
- Make a conscious effort to drink plenty of clear liquids before, during, and after taking your Suprep. You may drink water throughout the night and up to 4 hours prior to your procedure to stay hydrated.

THE DAY OF YOUR EXAM:

Remember not to have any food until after procedure!

- AT LEAST 4-5 HOURS BEFORE YOUR PROCEDURE START TIME, repeat steps 1-4 with the second 6 ounces Suprep bottle as described above. This may require you to get up in the middle of the night, if you have an early morning exam.
- You may have only clear liquids up until 4 hours prior to your procedure start time. We must insist on this "nothing by mouth" time for patient safety. Do not have anything by mouth after this point, as your stomach must be completely empty at the time of the procedure.
- If you have diabetes, test your blood sugar before coming in for your exam and bring the reading with you. If you take oral medications for your diabetes, don't take them the morning of your procedure unless otherwise instructed by your doctor. If you take insulin, take only ½ of your usual morning dose and bring the insulin with you to the procedure.
- You may take your usual medications (e.g. for blood pressure). Any medications taken the day of your procedure may be taken with a small sip of water, AT LEAST 4 HOURS PRIOR TO YOUR PROCEDURE START TIME.
- You may apply zinc oxide or petroleum jelly around the anus to prevent irritation from the prep. Baby wipes or adult diapers may be considered.

OTHER INFORMATION

- You must have someone available to drive you home after colonoscopy. If you are planning to take a taxi home, then a friend or family member must accompany you. Plan on spending 2-3 hours at the facility.
- You may receive up to 4 separate bills: 1-facility, 2-physician, 3-pathology, and 4-anesthesia.
- Wear comfortable clothing. Bring your glasses, hearing aids, inhaler, insurance cards, and completed information forms if applicable. Avoid bringing valuables. We will expect payment of co-pays, co-insurance, and deductibles at the time of service.
- If you must cancel or reschedule your procedure, then you must give notice 2 full business days in advance. Failure to do so will result in a \$100 cancellation fee.
- Additional information is available on our website: www.utahgastro.com